



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A23K 1/00, A23J 1/18, 1/20	A1	(11) International Publication Number: WO 00/35299 (43) International Publication Date: 22 June 2000 (22.06.00)
(21) International Application Number: PCT/NO99/00385 (22) International Filing Date: 16 December 1999 (16.12.99) (30) Priority Data: 19985919 17 December 1998 (17.12.98) NO (71) Applicant (for all designated States except US): SILDOLJE- OG SILDEMEILINDUSTRIENS FORSKNINGSINSTITUTT [NO/NO]; Kjerreidviken 16, N-5141 Fyllingsdalen (NO). (72) Inventors; and (75) Inventors/Applicants (for US only): SAMUELSEN, Tor, Andreas [NO/NO]; Vallalia 165, N-5050 Nesttun (NO). OTERHALS, Åge [NO/NO]; Geithusv. 76, N-5066 Hjellestad (NO). (74) Agent: AS BERGEN PATENTKONTOR; C. Sundtsgt. 36, N-5004 Bergen (NO).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: A POWDERED PROTEIN COMPOSITION HAVING BINDING PROPERTIES ADAPTED FOR USE AS PROTEIN SOURCE AND BINDER IN A FORMULATED FEED		
(57) Abstract A powdered protein composition which has a protein content of above 35 % by weight and which in formulated feed functions both as a protein source and as a binder comprises a matrix of protein containing particles of animal and/or vegetable origin and/or single cell protein, upon which a protein hydrolysate is dried. Production of the protein composition is described.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

5

A powdered protein composition having binding properties adapted for use as protein source and binder in a formulated feed.

10

The present invention relates to a powdered protein composition which has a protein content of above 35% and which in formulated feed functiones both as protein source and binder.

15

The invention also relates to a process for producing the protein composition.

Breeding of fish and other marine organisms increases to high extent, and the need for formulated feed is increasing. Compared to feed for ordinary domestic animals such a feed has a relatively high content of protein. An increasing part of the world's production of fish meal is used in formulated fish feed, in the form of pellets, and vegetable protein ingredients, single cell protein and other animal protein ingredients are incorporated as supplement. Earlier such feed contained considerable amounts of carbohydrates or other binders which provide a feed pellet having adequate strength, but which have low nutritive value. The development has proceeded towards a feed with a high energy level, and it is therefore less possibility for using binders having a low nutritive value, and problem arises because the feed pellet thus not have enough strength to resist the treatment it is exposed to during processing, storage and the transport.

20

25

30

One of the main ingredients in fish feed is fish meal, and the strength of the feed pellet is to a great extent determined by the physical properties of the fish meal,

35

CONFIRMATION COPY

which will vary with fish species, fishing season and the conditions during processing in the fish meal factory.

- Fish silage is in the literature first mentioned by Swedish researches (H. Edin, (1940): Nord. Jordbr. Forsk., Vol 22, p 142. - N. Olsson (1942): Landbrukshøgskolans Husdjursforsøksanstalt (1942), Report No. 7). Later a great number of articles have been published (Examples: I. Tatterson, in "Proc. of the Torry Res. Station Symp. on Fish Silage", Aberdeed (1976), p 1-10. - A. Gildberg, J. Raa (1979): Comp. Biochem. Physiol., 63B, p 309.- D. Potter, I. Tatterson, J. Wignall in "Advances in Fish Science and Technology" (ed. J.J. Connell), Fishing News Books Ltd. (1980), p 338-343).

- The fish silage is produced by grinding fish or parts of fish, and adding acid, usually formic acid, to a pH of 3.5 - 4. The ground fish can thereby be stored without bacterial decomposition, but the fish's own enzymes will cause a hydrolysis of the proteins to peptides and amino acids (S. Vecchi, Z. Coppes (1996): J. Food Biochem., Vol 20, p 193-214). Another, similar hydrolysate can be prepared in that proteinuous material, such as animal or vegetable protein or single cell protein, is treated with proteolytic enzymes in a manner known per se (J. Adler-Nissen "Enzymatic Hydrolysis of Food Proteins", Elsevier Applied Science Publishers, Ltd., 1986, G. Søbstad: "Meldinger fra SSF", No. 2, p 11-14, 1980 - F. Jacobsen, O. Lykke-Rasmussen: Process Biochem, 1984, p 165-169). The degree of hydrolysis can be determined in different ways, but is often defined as the percentage of protein (nitrogen x 6.25) which is not precipitated in water to which is added 5% trichloric acetic acid, i.e. that a degree of hydrolysis of 50% means that 50% of the protein is soluble in water which contains 5% trichloric acetic acid.

- Industrial processes are known where raw fish silage is heated and oil separated before the fish silage is concentrated by evaporation to a solid content of 30 -50% (B. Stormo, T. Strøm: "Ensilering av fiskeslo", FTFI-report no. 663.2-4-3, Fiskeriforskning, Tromsø, December 1978).

Both raw ensilage and fish silage concentrate are used as raw material for the production of feed for domestic animals and fish (F.E. Stone, (1989) Aquaculture, Vol 76 (1-2) p 109-118. - D. Manikandavelu, et al. (1992) Fish Technol., Vol 29 (2) p 111-113).

During the ensilaging process there will be a partial decomposition of some important amino acids (A. Gildberg, J. Raa: J. Sci. Food Agric., Vol 28, 1977, p 647). For animals with a high nutritive need, for example fish, the fish silage therefore should be used together with another valuable protein source, for example fish meal.

Protein hydrolysates have good binding properties in formulated feed and contribute in obtaining a feed pellet having the desired strength. With exception of the fish oil most of the raw materials which are used in such feed are present in the dry form, and for the process it would be an advantage if also the protein hydrolysate would be present as powder. Protein hydrolysate has a high content of ingredients with glue properties and will in dry form therefore appear as a hygroscopic powder with poor powder properties.

Mixing of such a powder into a feed blend is problematic.

An object of the present invention is to provide a protein composition which does not cause the problems mentioned above and which yield a feed pellet having adequate strength and a high energy content.

Further there is an object of the invention to provide a process for producing the protein composition.

These objects are obtained by the characterising features in claim 1 and claim 9, respectively.

Examples of matrix are press cake from the fish meal process, fish meal, meat/bone meal, soya meal, etc. It is thereby obtained protein compositions which in addition to good nutritive properties also have adequate powder technical properties, and, used in formulated feed, provide a pellet having adequate strength without use of other binders. Further, it has been shown that such protein

compositions can be used as binders during production of formulated feed wherein other raw materials yield poor binding properties.

The invention will be illustrated by means of the following examples with reference to the accompanying drawings, which show correlation between the amount of hydrolysate used and the properties of the composition which has been prepared.

10 Example 1

Feed pellets were prepared from two fish meals (F1 and F2), produced in the same factory from the same fish material, without and with addition of fish silage. F2 was produced by substituting 10% of the raw material to the cooker with fish silage. Dry feed mass introduced into the extruder contained 76% by weight of either F1 or F2. The feed mass also contained 19% by weight wheat meal. The fat content of the feed mass was adjusted to 11% by weight with fish oil before extrusion. Extrusion was carried out under similar conditions and with a 7 mm die opening. Feed pellets, A1 from feed mass containing F1 and A2 from feed mass containing F2, were dried to a water content of 7% by weight.

The following table shows comparison of physical quality of pellet from the two feed mixtures. Analysis of dried pellet was carried out before addition of oil to desired final level to obtain a higher fat level, for example up to 30 - 40%.

Feed pellet no.:	Pneumatic durability (%)	Hardness (kp)
A1	64	5.6
A2	80	9.7

30

Pneumatic durability is determined by a standardised method for simulated, pneumatic transport and state % of the part of the pellet which has not been crushed.

The figures for pneumatic durability show that the feed pellet from fish meal F2 has a higher strength and resist pneumatic transport better than the feed pellet from fish meal from F1.

5

Example 2

Feed pellets were prepared from two fish meals F3 and F4 containing different amounts of hydrolysed stick water protein as fish hydrolysate. The meals were produced in the same factory from the same fish raw material. F4 contained the double amount hydrolysed stick water protein compared to F3. The feed mass added to the extruder contained either F3 or F4. F3 contained 8.5% by weight hydrolysed protein, based on dry feed mass, and F4 contained 15.7% by weight. Otherwise the feed mass had the same content of other ingredients and was extruded/dried as described in example 1. Feed pellets produced, A3 from feed mass containing F3 and A4 from feed mass containing F4, were compared concerning physical quality. The analysis were carried out on dried pellet after addition of fish oil to desired final level.

Feed pellet no.:	Pneumatic durability (%)	Hardness (kp)
A3	0	1
A4	95	9.6

25

The figures show that feed pellet from F3 has a lower strength and does not resist pneumatic transport. Both strength and pneumatic durability are dramatically improved for feed pellet from F4.

30

Example 3

Based on fish raw material, namely stick water from the fish meal process, three fractions having different content of hydrolysed protein were produced:

P1: Stick water fraction containing 31% hydrolysed protein.

P2: Stick water fraction having 48% hydrolysed protein.

5 P3: Stick water fraction having 90% hydrolysed protein.

Starting with a fish meal (F5) and those three stick water fractions mentioned above 4 fish meals were prepared:

F5: Fish meal, 100%

10 F6: F5 containing P1 in the ratio 91.5% : 8.5%.

F7: F5 containing P2 in the ratio 91.5% : 8.5%.

F8: F5 containing P3 in the ratio 91.5% : 8.5%.

The feed mass introduced into the extruder contained either F5, F6, F7 or F8. F6 contained 9.1% of hydrolysed protein based on dried feed mass, F7 contained 9.7% by weight, and F8 contained 11% by weight. Otherwise the feed masses had an equal content of other ingredients and were extruded/dried as described in example 1. Feed pellets, A5, A6, A7 and A8 from feed masses containing F5, F6, F7 and F8 respectively, were compared what physical quality concerns. The analysis were carried out on dried pellet before addition of oil to desired final level.

25

Feed pellet no.:	Pneumatic durability (%)	Stability in water (%)
A5	0	74
A6	59	82
A7	69	84
A8	92	88

It appears from the table that the strength of the feed pellets, expressed by pneumatic durability, and the stability of the pellets in water increase with increasing addition of hydrolysed protein. If both pneumatic durability and stability in water were plotted as the

30

function of the amount of hydrolysed protein introduced into the extruder for A6, A7 and A8, respectively, a linear correlation as shown in figures 1 and 2, respectively, is obtained.

5

Example 4

A binder based on protein was prepared by drying fish silage on a matrix of fish meal. The binder (P4) contained 42% by weight fish silage, based on solid content. By
10 mixing fish meal (F9) and binder (P4) the following 5 fish meals were prepared:

F9: Fish meal, 100%.

F10: F9 containing P4 in the ratio 99.1% : 0.9%.

F11: F9 containing P4 in the ratio 98.3% : 1.7%.

15 F12: F9 containing P4 in the ratio 96.6% : 3.4%.

F13: F9 containing P4 in the ratio 91.4% : 8.6%.

The feed mass introduced into the extruder contained either F9, F10, F11, F12 or F13. Otherwise the feed mass
20 had the same content of other ingredients and was extruded/dried as described in example 1. The pellets, A9, A10, A11, A12 and A13 produced from feed mass containing F9, F10, F11, F12 and F13 respectively, were compared concerning physical quality. The analysis were carried out
25 on dried pellet before addition of oil to desired final level.

Feed pellet no.:	Mechanical durability (%)	Stability in water (%)
A9	52.2	77.5
A10	57.2	79.9
A11	60.8	80.2
A12	62.8	80.9
A13	80	84.4

30

Mechanical durability is determined according to a standardised method for simulating transport in bulk.

The durability of the pellets, expressed as mechanical durability, and the stability of the pellets in water increase with increasing addition of binder P4. If both mechanical durability and stability in water are plotted as the function of the amount of P4 introduced into the extruder for A9, A10, A11, A12 and A13 a linear correlation as shown in figures 3 and 4 is obtained.

10

C L A I M S.

1. Powdered protein composition which has a protein content of above 35% by weight and which in formulated feed functions both as a source of protein and as a binder, characterized in that it comprises a matrix of protein containing particles of animal and/or vegetable origin and/or single cell protein, upon which a protein hydrolysate has been dried.
2. Protein composition in accordance with claim 1, characterized in that the matrix comprises fish meal and/or press cake for preparing fish meal.
3. Protein composition in accordance with claim 1, characterized in that the matrix comprises meat/bone meal.
4. Protein composition in accordance with any of claims 1-3, characterized in that the matrix comprises meal of soya, raps, or other vegetable protein, or single cell protein, or mixtures thereof, having a protein content of at least 35% by weight.
5. Protein composition in accordance with any of the preceding claims, characterized in that the protein hydrolysate is prepared by hydrolysis of animal and/or vegetable protein and/or single cell protein.
6. Protein composition in accordance with claim 5, characterized in that the protein hydrolysate is fish silage, either in raw or concentrated form.

7. Protein composition in accordance any of the preceding claims, characterized in that the protein hydrolysate has a degree of hydrolysate of at least 50% by weight.

5

8. Protein composition in accordance any of the preceding claims, characterized in that the protein from the protein hydrolysate comprises 5 - 95% by weight of the solid content of the protein composition.

10

9. Process for preparing a powdered prote in composition which has a protein content of above 35% by weight and which in formulated feed functions both as a source of protein and as a binder, characterized
15 ed in that a protein hydrolysate in liquid form is dried on a matrix of particles of animal or vegetable origin and/or of single cell protein.

10. Process in accordance with claim 9, characterized
20 terized in that as a protein hydrolysate is used fish silage, hydrolysed fish protein, single cell protein, vegetable protein or a mixture thereof.

Figure 1.




Figure 2.




Figure 3.





Figure 4.



INTERNATIONAL SEARCH REPORT

International application No.

PCT/NO 99/00385

A. CLASSIFICATION OF SUBJECT MATTER

IPC7: A23K 1/00, A23J 1/18, A23J 1/20

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: A23K, A23J

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	File WPI, Derwent accession no. 1985-272846, AJINOMOTO GENERAL FOODS INC: "Improving desirability of dry and semi-dry pet foods - by incorporating crab metal and protein hydro- lysate"; & JP,A,60184360, 19850919 DW198544 --	1-10
X	GB 2247604 A (DIELEN LABORATORIES), 11 March 1992 (11.03.92) --	1-10
A	FR 2542013 A1 (BIO-INDUSTRIES), 7 Sept 1984 (07.09.84) --	1-10

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

29 March 2000

Date of mailing of the international search report

20-04-2000

Name and mailing address of the ISA/
Swedish Patent Office
Box 5055, S-102 42 STOCKHOLM
Facsimile No. +46 8 666 02 86

Authorized officer

Eva Johansson/EÖ
Telephone No. +46 8 782 25 00

Form PCT/ISA/210 (second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/NO 99/00385

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 3697285 A (WILLIAM T. FAITH), 10 October 1972 (10.10.72) --	1-10
A	US 4764383 A (SHERWIN R. BROWN ET AL), 16 August 1988 (16.08.88) --	1-10
A	WO 8704051 A1 (PETROFF, PAUL), 16 July 1987 (16.07.87) -- -----	1-10

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT
Information on patent family members

02/12/99

International application No.

PCT/NO 99/00385

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB 2247604 A	11/03/92	ES 2035781 B FR 2663200 A,B IT 1249663 B IT T0910462 D PT 98037 A,B	16/02/94 20/12/91 09/03/95 00/00/00 30/04/92
FR 2542013 A1	07/09/84	NONE	
US 3697285 A	10/10/72	BR 6915318 D NL 7014806 A	00/00/00 14/04/71
US 4764383 A	16/08/88	NONE	
WO 8704051 A1	16/07/87	DK 474187 A FI 860144 D FI 860623 A JP 63502159 T NO 873814 A	11/09/87 00/00/00 14/07/87 25/08/88 11/09/87

Form PCT/ISA/210 (patent family annex) (July 1992)

THIS PAGE BLANK (USPTO)